

**Question:** What gas mixtures are best for modified atmosphere packaging (MAP) to store ready-to-eat (RTE) meat products to obtain a 2-week shelf-life and not impact quality/flavor/color?

**Response:** Most studies that compared different gas mixtures in MAP during storage of meat products looked at fresh meat, and those that did look at RTE meats were usually more interested in safety than quality. In general, studies in the literature found that a high (at least 30%) CO<sub>2</sub> concentration and an absence of O<sub>2</sub> seem to be the preferred MAP conditions for RTE meat products. Here are some of the key studies identified:

| Reference                 | Meat product                                      | MAP gas composition                                                                                                                                      | Storage conditions | Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|---------------------------|---------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (Stasiewicz et al., 2015) | Pre-cooked sausages in artificial protein casings | 20% CO <sub>2</sub> + 80% N <sub>2</sub><br>50% CO <sub>2</sub> + 50% N <sub>2</sub><br>80% CO <sub>2</sub> + 20% N <sub>2</sub>                         | 4°C for 15 days    | <ul style="list-style-type: none"> <li>The presence of carbon dioxide in modified atmosphere slowed down the rate of lipid oxidation in meat products.</li> <li>The atmosphere with the highest carbon dioxide concentrations was most suitable for storing pre-cooked sausages.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| (Gokoglu et al., 2010)    | Beef frankfurter-type sausages                    | 30% CO <sub>2</sub> + 70% N <sub>2</sub><br>70% CO <sub>2</sub> + 30% N <sub>2</sub><br>100% CO <sub>2</sub><br>80% CO <sub>2</sub> + 20% O <sub>2</sub> | 4°C for 28 days    | <ul style="list-style-type: none"> <li>CO<sub>2</sub> was effective at both inhibiting oxidation and in microbial inhibition.</li> <li>The highest viable count was found in the 30% CO<sub>2</sub> + 70% N<sub>2</sub> atmosphere; the lowest with 100% CO<sub>2</sub>.</li> <li>The lowest count for lactic acid bacteria was found for the 80% CO<sub>2</sub> + 20% O<sub>2</sub> atmosphere, but this atmosphere had the lowest sensory score.</li> <li>The authors concluded the best atmosphere to maximize quality and shelf-life was 70% CO<sub>2</sub> + 30% N<sub>2</sub>.</li> </ul>                                                                                                                                                                    |
| (Summo et al., 2016)      | Ripened sausages containing 120 mg/kg nitrate     | 70% N <sub>2</sub> + 30% CO <sub>2</sub><br>80% N <sub>2</sub> + 20% CO <sub>2</sub><br>95% N <sub>2</sub> + 5% CO <sub>2</sub>                          | 4°C for 5 months   | <ul style="list-style-type: none"> <li>Oxidative degradation occurred less at higher CO<sub>2</sub> levels (probably during the initial storage period when residual oxygen from the food product was still available).</li> <li>Sensory scores were better at higher CO<sub>2</sub> levels.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| (Cachaldora et al., 2013) | Morcilla (cooked blood sausage)                   | 15% O <sub>2</sub> + 35% N <sub>2</sub> + 50% CO <sub>2</sub><br>60% N <sub>2</sub> + 40% CO <sub>2</sub><br>40% N <sub>2</sub> + 60% CO <sub>2</sub>    | 4°C for 8 weeks    | <ul style="list-style-type: none"> <li>Packaging did not influence shelf life, which was greater than 8 weeks for all packaging conditions.</li> <li>Samples packaged with high CO<sub>2</sub> (40% N<sub>2</sub> plus 60% CO<sub>2</sub>) should the lowest values of TBARS (a sign of lipid oxidation) at the end of storage</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                          |
| (Jin and Choi, 2017)      | Fermented dry-cured hams                          | Aerobic packaging<br>Vacuum packaging<br>30% CO <sub>2</sub> + 70% N <sub>2</sub>                                                                        | 4°C for 12 weeks   | <ul style="list-style-type: none"> <li>In terms of CIE*color, TBARS, VBN, and sensory evaluation, the vacuum packaging and MAP methods better maintained the quality characteristics of fermented dry-cured hams during storage when compared with aerobic packaging.</li> <li>In comparisons between the vacuum packaging and MAP methods, the values of pH, moisture and tenderness were higher, and that of salt content was lower in vacuum packaging.</li> <li>Additionally, the redness value, which is regarded as a color stability criterion, and inhibition of microorganism growth were higher at the end of the storage periods for vacuum packaging vs. MAP.</li> <li>The VBN value was lower for the MAP packaging method during storage.</li> </ul> |
| (Naas et al., 2013)       | Low-fat turkey bologna                            | 100% CO <sub>2</sub><br>Air<br>Vacuum                                                                                                                    | 4°C for 42 days    | <ul style="list-style-type: none"> <li>100% CO<sub>2</sub> packaging prevented outgrowth of <i>Listeria monocytogenes</i> throughout 42 d of storage, whereas non- CO<sub>2</sub> packaging displayed a 2-log increase in population during storage.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |

| Reference             | Meat product                          | MAP gas composition                                             | Storage conditions                | Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|-----------------------|---------------------------------------|-----------------------------------------------------------------|-----------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (Smiddy et al., 2002) | Various sliced processed cooked meats | 70% N <sub>2</sub> and 30% CO <sub>2</sub> ;                    | 4°C for 21 days                   | <ul style="list-style-type: none"> <li>Although all products (25 packs each of 13 different cooked meat products from 3 different manufacturers) were all packaged with the same MAP, O<sub>2</sub> was present in 88% of packages within 24 hours of packaging (20% were &gt;1.2% O<sub>2</sub>). By day 21, 79.3% of packages contained &gt;1.2% O<sub>2</sub>.</li> </ul>                                                                                                                                                                                                                                                                                                         |
| (Lawlor et al., 2000) | Cooked, uncured turkey                | 100% N <sub>2</sub><br>30% CO <sub>2</sub> + 70% N <sub>2</sub> | 4, 10, and 15°C for up to 60 days | <ul style="list-style-type: none"> <li>Packaging atmosphere had no significant effect on APC.</li> <li>Total aerobic populations of inoculated turkey packaged under 30% CO<sub>2</sub>:70% N<sub>2</sub> were 1 to 2.5 logs lower than in inoculated turkey packaged under 100% N<sub>2</sub>, an effect that became more pronounced as storage temperature decreased.</li> <li>LAB counts were not affected by packaging atmosphere at any temperature.</li> <li>The number of days until sensory unacceptability occurred was in some cases the same for the two gas compositions; no consistent influence of gas composition on sensory unacceptability was apparent.</li> </ul> |

### References:

- Cachaldora A, Garcia G, Lorenzo JM, Garcia-Fontan MC (2013) Effect of modified atmosphere and vacuum packaging on some quality characteristics and the shelf-life of "morcilla", a typical cooked blood sausage. *Meat Sci* 93 (2):220-225.
- Gokoglu N, Yerlikaya P, Uran H, Topuz OK (2010) THE EFFECT OF MODIFIED ATMOSPHERE PACKAGING ON THE QUALITY AND SHELF LIFE OF FRANKFURTER TYPE-SAUSAGES. *J Food Qual* 33:367-380.
- Jin SK, Choi JS (2017) Effect of packaging methods on quality characteristics of fermented dry-cured hams during cold storage at 4 A degrees C. *J Food Meas Charact* 11 (4):2052-2060.
- Lawlor KA, Pierson MD, Hackney CR, Claus JR, Marcy JE (2000) Nonproteolytic *Clostridium botulinum* toxigenesis in cooked turkey stored under modified atmospheres. *J Food Prot* 63 (11):1511-1516.
- Naas H, Martinez-Dawson R, Han I, Dawson P (2013) Effect of combining nisin with modified atmosphere packaging on inhibition of *Listeria monocytogenes* in ready-to-eat turkey bologna. *Poult Sci* 92 (7):1930-1935.
- Smiddy M, Papkovsky D, Kerry J (2002) Evaluation of oxygen content in commercial modified atmosphere packs (MAP) of processed cooked meats. *Food Res Int* 35 (6):571-575.
- Stasiewicz M, Lipinski K, Cierach M, Purwin C (2015) Lipid oxidation in meat products packaged stored under vacuum and modified atmosphere with different concentrations of carbon dioxide and nitrogen. *J Food Saf Food Qual* 66 (5):136-140.
- Summo C, Pasqualone A, Paradiso V, Centomani I, Centoducati G, Caponio F (2016) Lipid degradation and sensory characteristics of ripened sausages packed in modified atmosphere at different carbon dioxide concentrations. *J Sci Food Agr* 96 (1):262-270.

**Compiled by W. Bedale, Food Research Institute, University of Wisconsin-Madison; [bedale@wisc.edu](mailto:bedale@wisc.edu)**